

AD-A257 652



2

TGAL-92-04

HIGH-PRECISION RELATIVE EVENT LOCATION WITH CROSS-SPECTRAL ANALYSIS

Indra N. Gupta
Peter Davis
Thomas W. McElfresh
Robert A. Wagner

Teledyne Geotech Alexandria Laboratories
314 Montgomery Street
Alexandria, VA 22314-1581

DTIC
ELECTE
NOV 13 1992
S C D

APRIL 1992

SEMI-ANNUAL REPORT (23 August 1991 - 4 April 1992)

Approved for Public Release; Distribution Unlimited

Prepared for:
UNITED STATES AIR FORCE
AIR FORCE SYSTEM COMMAND
PHILLIPS LABORATORY (PL)
KIRTLAND AFB, NM 87117-6008

Monitored by:
DEFENSE ADVANCED RESEARCH PROJECTS AGENCY
NUCLEAR MONITORING RESEARCH OFFICE
3701 NORTH FAIRFAX DRIVE
ARLINGTON, VA 22203-1714

The views and conclusions contained in this report are those of the authors and should not be interpreted as representing the official policies, either expressed or implied, of the Defense Advanced Research Projects Agency or the U.S. Government.

40-257
92-29353



3388

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 4 April 1992		3. REPORT TYPE AND DATES COVERED Technical Report, 23 Aug 1991 - 4 Apr 1992
4. TITLE AND SUBTITLE High-Precision Relative Location with Cross-Spectral Analysis			5. FUNDING NUMBERS Contract F29601-91-C-DB07	
6. AUTHOR(S) I. N. Gupta, J. P. Davis, T. W. McElfresh, and R. A. Wagner				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Teledyne Geotech 314 Montgomery Street Alexandria, VA 22314-1581			8. PERFORMING ORGANIZATION REPORT NUMBER TGAL-92-04	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) DARPA-NMRO 3701 North Fairfax Drive, #717 Arlington, VA 22203-1714			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) <p>Location relative to a reference event is often more useful and precise than absolute event location. The use of relative location accounts for most of the errors arising from path effects. Such location requires relative times for pairs of events observed at a common station, which can be obtained with high precision for similar events by waveform cross-correlation. The precision can be further boosted by applying the cross-spectral analysis method, which can obtain relative delay times with resolution up to an order magnitude better than the seismogram sampling interval. During this report period, records of a large number of explosions with precisely known locations were assembled. Software for determining the delay time between two waveforms was developed and tested on real data. Preliminary analysis of data from six closely-located Yucca Flat explosions recorded at the four broadband digital stations, ELK, KNB, LAC, and MNV provided encouraging results. The mean location error was found to be only about 1 km; an impressive result if one considers the large epicentral distances (about 200 km to 320 km) and the complex geology of the Nevada Test Site.</p>				
14. SUBJECT TERMS Relative Location, Cross-spectral Analysis, Waveform Cross-correlation			15. NUMBER OF PAGES 37	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	

1. SUMMARY

Location relative to a reference event is often more useful and precise than absolute event location. The use of relative location accounts for most of the errors arising from path effects. Such location requires relative times for pairs of events observed at a common station, which can be obtained with high precision for similar events by waveform cross-correlation.

The precision can be further boosted by applying the cross-spectral analysis method, which can obtain relative delay times with resolution up to an order of magnitude better than the seismogram sampling interval. In the cross-spectral analysis method, the delay time is simply the slope of the phase of the cross-spectrum. This is obtained by fitting a slope through 0, using a weighting scheme based on the coherence of the cross-spectrum.

During this report period, records of a large number of explosions with precisely known locations were assembled. Software for determining the delay time between two waveforms was developed and tested on real data. Preliminary analysis of data from six closely-located Yucca Flat explosions recorded at the four broadband digital stations, ELK, KNB, LAC, and MNV provided encouraging results. The relative arrival times between two closely spaced explosions recorded at a common station were determined with a precision of 0.001 sec. The mean location error was found to be only about 1 km; an impressive result if one considers the large epicentral distances (about 200 km to 320 km) and the complex geology of the Nevada Test Site.

DTIC QUALITY INSPECTED 4

Accession For	
NTIS	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

(THIS PAGE INTENTIONALLY LEFT BLANK)

TABLE OF CONTENTS

	Page
1. SUMMARY	iii
2. INTRODUCTION	1
3. CROSS-SPECTRAL ANALYSIS	3
4. PRELIMINARY RESULTS	5
5. CONCLUSIONS	19
6. ACKNOWLEDGMENTS	20
7. REFERENCES	21
DISTRIBUTION LIST	23

(THIS PAGE INTENTIONALLY LEFT BLANK)

2. INTRODUCTION

One of the most important tasks in monitoring underground nuclear tests is the simple location of regional events. In fact, the location itself is often useful for discriminating between explosions and earthquakes. Location techniques generally rely on travel times and back azimuths recorded at several stations, using some variant on Geiger's classic inversion algorithm (*e. g.*, Aki and Richards, 1980). The main cause of errors in such location algorithms is poorly known earth structure models or three-dimensional variations in velocity. While these effects tend to be averaged out if enough stations are used, locations using small numbers of stations can have large errors. However, in many cases, a location relative to a reference event is sufficient. The most common use of relative location in seismology is locating aftershocks relative to the main shock. This is useful because the smaller events are observed at fewer stations and thus their locations are affected more by earth structure. Relative relocation has the advantage of canceling out the major portion of the path effect, if the event is close enough to the reference event.

Another case where relative location can be useful is in locating an event whose absolute location is near a known mine or quarry, or near a known nuclear testing site. In this case, the probability that a given event occurred within a mine or test site could be established with greater certainty than with an absolute location by locating the event relative to a reference event known to have occurred at the mine or test site. Relative location can also improve the locations of nuclear tests, particularly if the absolute location of a reference event can be established using satellite imagery.

In conventional relative event location, events are located relative to a reference event by using time differences for a given arrival measured for both the reference event and the other

events. Lynnes and Ruff (1985) made a comparison of the ISC spatial locations of an aftershock sequence for an earthquake in the North Atlantic and relative locations based only on differences between aftershock and main shock P wave arrival times. Despite the fact that many of the events were recorded by as few as three stations, the relative location method produced locations aligned along the fault plane.

3. CROSS-SPECTRAL ANALYSIS

Relative locations are limited primarily by the precision with which the phase arrival times can be measured. For two events that occur close together, the waveforms are often similar. In such cases a cross-correlation of an event's waveforms can provide much improved estimates of the relative times, since the full waveform is used rather than a subjective pick of the initial phase arrival. Conventional cross-correlation slides one waveform past another, and the lag with the optimum correlation coefficient is taken as the relative time. However, the cross-spectral analysis method can yield improved relative time estimates. Following Ito (1985), the cross-spectrum, $Sc(f)$, of two signals A and B is:

$$Sc(f) = \frac{F_a(f) F_b^*(f)}{T} = \text{Re}(f) - i \text{Im}(f), \quad (1)$$

where $F(f)$ is the Fourier transform of $f(t)$ and $*$ indicates the complex conjugate. Then the phase spectrum $\phi(f)$ is

$$\phi(f) = \tan^{-1} \frac{-\text{Im}(f)}{\text{Re}(f)} \quad (2)$$

If the two seismograms are similar in shape but have different amplitudes, then

$$f_b(t) = k f_a(t+\tau), \quad (3)$$

and by the application of the shift theorem, the phase spectrum becomes

$$\phi(f) = 2 \pi f \tau \quad (4)$$

Thus the delay time τ can be obtained by simply by fitting a straight line through the phase of the cross-spectrum with intercept 0. In fitting this slope, the values are weighted based on the coherence $C(f)$:

$$C(f) = \frac{\left| \sum_{f'=f-\Delta f}^{f'=f+\Delta f} S_c(f') \right|}{\sum_{f'=f-\Delta f}^{f'=f+\Delta f} \sqrt{S_a(f') S_b(f')}} \quad (5)$$

where $S(f)$ is the signal power. The weighting factor, $W(f)$ in the linear fit is based on the Hannon-Thomson processor (Knapp and Carter, 1976), *i. e.*



$$W(f) = \frac{C^2(f)}{1 - C^2(f)} \quad (6)$$

The slope of the line has a continuum of possible values, rather than discrete values, with the result that estimates of delay time for highly coherent pairs can actually be an order of magnitude more precise than the sample rate. This technique has been applied successfully to local sequences of earthquakes with impressively precise locations. Fremont and Malone (1987) obtained precisions of 20 m for earthquakes within 250 m of each other. Ito (1985) obtained precision of 50 m for microearthquakes within about 500 m of each other.

4. PRELIMINARY RESULTS

In order to understand the influence of path differences on waveforms, records of a large number of explosions with precisely known locations have been assembled. Most of these data are at regional distances and the magnitude, m_b , varies over a large range (about 3.3 to 6.2). An analysis of differences in the observed regional phases (especially Pn and Lg) from closely spaced explosions at the Nevada Test Site recorded at the four broadband digital stations, ELK, KNB, LAC, and MNV has been started. Attempts will be made to understand the observed spectral and time-domain differences in terms of variations of inter-shot distance, shot depth, and geological environment.


Software for determining the delay time between two waveforms has been developed and tested. The relative arrival times between two events recorded at a common station are determined with a precision of 0.001 sec. Results based on use of the Yucca Flat explosions ROUSANNE and BASEBALL, assuming BASEBALL to be the shot with known location, derived from records at ELK, KNB, LAC, and MNV are shown in Figures 1, 2, 3, and 4, respectively. The geographic locations of these two shots, separated by about 2.37 km, are precisely known. The digital data are sampled at about 42 samples/sec. Figure 1 (bottom) shows the input waveforms which may be selected to be of any desired duration. The signal window over which a cosine taper applies is also variable. The top plot shows the phase of the cross-spectrum varying between -2π and $+2\pi$. The lower plot shows the coherency, varying between 0 and 1, derived by using a bandwidth of 1.0 Hz. As expected from theory (*e. g.*, Frankel and Clayton, 1986), the coherency is generally observed to decline with increasing frequency. The delay time is simply the slope of the phase of the cross-spectrum over the specified range of 0.5-5.0 Hz and is obtained by fitting a slope through 0, using a



1177.9 nm 0-P Seis 1 ELK BASEBALLelk
 Start time: 15 Jan 81 20:25:57.3182

256 pts
 42.010 samp/sec
 Cosine Taper
 Backup: 42 pts


Start Point
 7657



979.3 nm 0-P Seis 1 ELK ROUSANNEelk
 Start time: 12 Nov 81 15:0:57.2112

Corrected Arrival:
 15:0:58.2079

Start Point
 7652



0.0 5.0 Sec

Figure 1. Cross-spectral analysis for measuring delay times. The two input waveforms (bottom), each 256 points long, provide the phase of the cross-spectrum (top) and coherence based on bandwidth of 1.0 Hz for frequencies up to 10 Hz. The slope of the phase spectrum, computed by fixing the intercept to 0 and applying weights related to the coherence, provides the delay time used to obtain the corrected arrival time for ROUSANNE recorded at ELK.









Figure 2. Similar to Figure 1 for records at KNB.



1747.3 nm 0-P Seis 1 LAC BASEBALLlacv
 Start time: 15 Jan 81 20:25:43.5597

256 pts
 42.010 samp/sec
 Cosine Taper
 Backup: 42 pts



967.0 nm 0-P Seis 1 LAC ROUSANNElacv
 Start time: 12 Nov 81 15:0:43.8335

Corrected Arrival:
 15:0:44.8394




Figure 3. Similar to Figure 1 for records at LAC.






Figure 4. Similar to Figure 1 for records at MNV.

weighting scheme based on coherence of the cross-spectrum (see equation 6).

Using BASEBALL as the reference explosion, the computed arrival times (such as those indicated on Figures 1 to 4) are used as input to the LOCATE feature of the Analyst Review Station (ARS) to obtain the epicentral location and the origin time. The arrival times (without any delay correction) for BASEBALL recorded at the four stations are also used to compute the corresponding ARS location. The ARS computed locations and the actual (known) locations are given in Table 1 and also shown in Figure 5. The ARS computed locations for the five shots (paired with BASEBALL) are shifted by the amount of shift between the ARS computed and actual locations of BASEBALL. A comparison of the shifted locations of each of the five shots with its actual location provides the location error associated with each shot (Figure 5). The mean location error is only 1.22 km with one standard deviation of 0.61 km. This result appears to be impressive if one considers the large epicentral distances (about 200 km to 320 km) and the complex geology of the Nevada test site.

Reference: BASEBALL
256 pt window







Figure 5. Map of six explosions at Yucca Flat showing the actual locations, ARS computed locations, and locations with shift based on BASEBALL. The five lines indicate the error between the actual and computed locations.

TABLE 1


CROSS-SPECTRAL ANALYSIS (SIX YUCCA FLAT SHOTS)

EVENT	ACTUAL LOCATION	COMPUTED LOCATION	SHIFTED LOCATION	SHIFTED - ACTUAL	LOCATION ERROR (km)
BASEBALL	37.0871	37.0769	37.0871	0.0000	0.000
	-116.0447	-116.0631	-116.0447	0.0000	
DAUPHIN	37.1115	37.0951	37.1053	-0.0062	1.152
	-116.0187	-116.0475	-116.0291	-0.0104	
HEARTS	37.0881	37.0749	37.0851	-0.0030	0.818
	-116.0528	-116.0628	-116.0444	0.0084	
PALIZA	37.0816	37.0523	37.0625	-0.0191	2.215
	-116.0088	-116.0344	-116.0160	-0.0072	
ROUSANNE	37.1082	37.0924	37.1026	-0.0056	0.634
	-116.0490	-116.0660	-116.0476	0.0014	
TILCI	37.0763	37.0564	37.0666	-0.0097	1.291
	-116.0685	-116.0949	-116.0765	-0.0080	

Cross spectral analysis based on the use of shorter (128 points) windows was also carried out for the same 6 explosions. Results for ROUSANNE and BASEBALL (reference shot) are shown in Figure 6, 7, 8, and 9 in which a bandwidth of 2.0 Hz is used for deriving coherency and a frequency range of 1.0-10.0 Hz is used for computing the delay time. A comparison with the earlier results (Figures 1 to 4) suggests significantly larger coherency over a much larger range of frequencies. It seems therefore that the first arrivals in Pn are considerably more similar than the later arriving phases, perhaps due to greater contamination by later-arriving scattered energy. As before, the coherency is generally observed to decline with





128 pts
 42.010 samp/sec
 Cosine Taper
 Backup: 21 pts



Corrected Arrival:
 15 : 0 : 58.1956

Figure 6. Similar to Figure 1 for shorter (128 points) waveforms recorded at ELK.



3972.5 nm 0-P Seis 1 KNB BASEBALLknbv
 Start time: 15 Jan 81 20:25:42.6075

128 pts
 42.010 samp/sec
 Cosine Taper
 Backup: 21 pts





2289.4 nm 0-P Seis 1 KNB ROUSANNEknbv
 Start time: 12 Nov 81 15:0:42.6671

Corrected Arrival:
 15:0:43.1706




0.0 5.0 Sec

Figure 7. Similar to Figure 1 for shorter (128 points) waveforms recorded at KNB.




1788.8 nm 0-P Seis 1 LAC BASEBALLlacv
Start time: 15 Jan 81 20:25:44.0595



128 pts
42.010 samp/sec
Cosine Taper
Backup: 21 pts

1014.1 nm 0-P Seis 1 LAC ROUSANNElacv
Start time: 12 Nov 81 15:0:44.3096



Corrected Arrival:
15:0:44.8144

0.0 5.0 Sec

Figure 8. Similar to Figure 1 for shorter (128 points) waveforms recorded at LAC.







Figure 9. Similar to Figure 1 for shorter (128 points) waveforms recorded at MNV.

increasing frequency. The location results, shown in Figure 10, are very similar to those in Figure 5 in both the magnitude and azimuthal direction of the location errors. The mean location error is 1.31 km with a standard deviation of 0.62 km.

A limitation of our analysis so far has been the use of the ARS to derive absolute locations for the reference and other shots. In the future, we plan to obtain relative locations directly from differences between arrival times of the reference shot and other explosions. This may significantly improve the precision of relative locations.

Reference: BASEBALL
128 pt window



B BASEBALL	* ACTUAL
R ROUSANNE	O ARS COMPUTED
H HEARTS	X SHIFTED
D DAUPHIN	
P PALIZA	
T TILCI	

Figure 10. Similar to Figure 5 based on the use of shorter (128 points) waveforms. Note the similarity of location errors with those in Figure 5.

5. CONCLUSIONS

Good quality records of a large number of explosions with precisely known locations and reliable time channel were assembled and software for determining the delay time between two waveforms was developed and tested. Encouraging results were obtained in a preliminary analysis of data from six closely-located Yucca Flat explosions recorded at the four broadband digital stations, ELK, KNB, LAC, and MNV. The relative arrival times between two closely spaced explosions recorded at a common station were determined with a precision of 0.001 sec. Use of long and short signal windows gave nearly identical results with the mean location error of only about 1 km. This is an impressive result if one considers the large epicentral distances (about 200 km to 320 km) and the complex geology of the Nevada test site.

6. ACKNOWLEDGMENTS

We thank Chris Lynnes for valuable advice and suggestions during the course of this study. This research was funded by the Nuclear Monitoring Research Office of the Defense Advanced Research Projects Agency under Contract F29601-91-C-DE07. The views and conclusions contained in this paper are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the Defense Advanced Research Projects Agency or the U. S. Government.

7. REFERENCES

- Aki, K., and P. G. Richards (1980). *Quantitative Seismology, Theory and Methods*, vol. 2, W. H. Freeman and Co., San Francisco.
- Frankel, A. and R. W. Clayton (1986). Finite difference simulations of seismic scattering: implications for the propagation of short-period seismic waves in the crust and models of crustal heterogeneity, *J. Geophys. Res.* 91, 6465-6489.
- Fremont, M. J. and S. D. Malone (1987). High precision relative locations of earthquakes at Mount St. Helens, Washington, *J. Geophys. Res.* 92, 10223-10236.
- Ito, A. (1985). High resolution relative hypocenters of similar earthquakes by cross-spectral analysis method, *J. Phys. Earth* 33, 279-294.
- Knapp, C. H., and G. C. Carter (1976). The generalized method for estimation of time delay, *IEEE Transactions on Acoustics, Speech, and Signal Processing* 24, 320-327.
- Lynnes, C. S., and L. J. Ruff (1985). Source process and tectonic implications fo the great 1975 North Atlantic earthquake, *Geophys. J. R. astr. Soc.* 82, 497-510.

(THIS PAGE INTENTIONALLY LEFT BLANK)

NON-GOVERNMENT CONTRACTORS

Prof. Thomas Ahrens
Seismological Lab, 252-21
Div. of Geol. & Planetary Sciences
California Institute of Technology
Pasadena, CA 91125

Dr. Thomas C. Bahe, Jr.
Dr. Thomas J. Serena, Jr.
Science Applications Int'l Corp.
10260 Campus Point Drive
San Diego, CA 92121
(2 copies)

Dr. Peter Basham
Dr. Robert North
Earth Physics Branch
Geological Survey of Canada
1 Observatory Crescent
Ottawa, Ontario, CANADA K1A 0Y3

Dr. Douglas R. Baumgardt
Dr. Zoltan Daz
ENSCO, Inc.
5400 Port Royal Road
Springfield, VA 22151-2388

Prof. Jonathan Berger
IGPP, A-025
Scripps Institution of Oceanography
University of California, San Diego
La Jolla, CA 92093

Dr. G. A. Bollinger
Department of Geological Sciences
Virginia Polytechnic Institute
21044 Derring Hall
Blacksburg, VA 24061

The Librarian
Dr. Jerry Carter
Dr. Stephen Bratt
Center for Seismic Studies
1300 North 17th Street, Suite 1450
Arlington, VA 22209-2308
(3 copies)

Michael Browne
Teledyne Geotech
3401 Shiloh Road
Garland, TX 75041

Dr. Lawrence J. Burdick
Woodward-Clyde Consultants
566 El Dorado Street
Pasadena, CA 91109-3245

Dr. Theodore Cherry
Science Horizons, Inc.
710 Encinitas Blvd., Suite 200
Encinitas, CA 92024 (2 copies)

Dr. Kin Yip Chun
Geophysics Division
Physics Department
University of Toronto
Ontario, CANADA M5S 1A7

Dr. Paul M. Davis
Dept. Earth & Space Sciences
University of California (UCLA)
Los Angeles, CA 90024

Prof. Steven Day
Department of Geological Sciences
San Diego State University
San Diego, CA 92182

Ms. Eva Johannisson
Senior Research Officer
National Defense Research Institute
P.O. Box 27322
S-102 54 Stockholm, SWEDEN

Dr. Mark D. Fisk
Mission Research Corporation
735 State Street
P.O. Drawer 719
Santa Barbara, CA 93102

Prof. Stanley Flatte
Applied Sciences Building
University of California
Santa Cruz, CA 95064

Robert C. Kemerait
ENSCO, Inc.
445 Pineda Court
Melbourne, FL 32940

Dr. Roger Fritzel
Pacific Sierra Research
1401 Wilson Blvd., Suite 1100
Arlington, VA 22209

Prof. Brian L. N. Kennett
Research School of Earth Sciences
Institute of Advanced Studies
G.P.O. Box 4
Canberra 2601, AUSTRALIA

Dr. Holly K. Given
Inst. Geophys. & Planet. Phys.
Scripps Inst. Oceanography (A-025)
University of California-San Diego
La Jolla, CA 92093

Dr. Richard LaCoss
MIT-Lincoln Laboratory
M-200B
P.O. Box 73
Lexington, MA 02173-0073

Prof. Hans-Peter Harjes
Institute for Geophysik
Ruhr University/Bochum
P.O. Box 102148
4630 Bochum 1, FRG

Prof. Fred K. Lamb
Univ. of Illinois
Department of Physics
1110 West Green Street
Urbana, IL 61801

Prof. Donald V. Helmberger
Seismological Laboratory
Div. of Geol. & Planetary Sciences
California Institute of Technology
Pasadena, CA 91125

Prof. Charles A. Langston
Geosciences Department
403 Deike Building
The Pennsylvania State University
University Park, PA 16802

Prof. Eugene Herrin
Prof. Brian Stump
Inst. for the Study of Earth and Man
Geophysical Laboratory
Southern Methodist University
Dallas, TX 75275

Prof. Thorne Lay
Dr. Susan Schwartz
Institute of Tectonics
Earth Science Board
University of California, Santa Cruz
Santa Cruz, CA 95064

Prof. Bryan Isacks
Prof. Muawia Barazangi
Cornell University
Department of Geological Sciences
SNEE Hall
Ithaca, NY 14850

Prof. Arthur Lerner-Lam
Prof. Paul Richards
Prof. C. H. Scholz
Lamont-Doherty Geol. Observatory
of Columbia University
Palisades, NY 10964

Prof. Lane R. Johnson
Prof. Thomas V. McEvilly
Seismographic Station
University of California
Berkeley, CA 94720

Dr. Manfred Henger
Fed. Inst. for Geosci. & Nat'l Res.
Postfach 510153
D-3000 Hanover 51, FRG

Dr. Peter Marshall
Procurement Executive
Ministry of Defense
Blacknest, Brimpton
Reading FG7-4RS, UNITED KINGDOM

Dr. Randolph Martin, III
New England Research, Inc.
76 Olcott Drive
White River Junction, VT 05001

Dr. Bernard Massinon
Societe Radiomana
27 rue Claude Bernard
75005 Paris, FRANCE (2 copies)

Dr. Gary McCartor
Prof. Henry L. Gray
Department of Physics
Southern Methodist University
Dallas, TX 75275

Dr. Keith L. McLaughlin
S-CUBED
P.O. Box 1620
La Jolla, CA 92038-1620

Dr. Pierre Mecheler
Societe Radiomana
27 rue Claude Bernard
75005 Paris, FRANCE

Prof. Bernard Minster
Prof. John Orcutt
Dr. Holly Given
IGPP, A-025
Scripps Institute of Oceanography
University of California, San Diego
La Jolla, CA 92093

Prof. Brian J. Mitchell
Dr. Robert Herrmann
Dept of Earth & Atmospheric Sciences
St. Louis University
St. Louis, MO 63156

Mr. Jack Murphy
S-CUBED
11800 Sunrise Valley Drive
Suite 1212
Reston, VA 22091
(2 copies)

Dr. Jay J. Pulli
Radix Systems, Inc.
2 Taft Court, Suite 203
Rockville, MD 20850

Dr. Frode Ringdal
Dr. Svein Mykkeltveit
NTNF/NORSAR
P.O. Box 51
N-2007 Kjeller, NORWAY
(2 copies)

Dr. Wilmer Rivers
Teledyne Geotech
314 Montgomery Street
Alexandria, VA 22314
(2 copies)

Dr. Richard Sailor
TASC, Inc.
55 Walkers Brook Drive
Reading, MA 01867

Prof. Charles G. Sammis
Prof. Kei Aki
Center for Earth Sciences
University of Southern California
University Park
Los Angeles, CA 90089-0741

Prof. David G. Simpson
Lamont-Doherty Geological Observatory
of Columbia University
Palisades, NY 10964

Dr. Stewart W. Smith
Geophysics AK-50
University of Washington
Seattle, WA 98195

Prof. Clifford Thurber
Prof. Robert P. Meyer
University of Wisconsin-Madison
Department of Geology & Geophysics
1215 West Dayton Street
Madison, WI 53706

Prof. K. Nafi Toksoz
Prof. Anton Dainty
Earth Resources Lab
Mass. Institute of Technology
42 Carleton Street
Cambridge, MA 02142

Prof. Terry C. Wallace
Department of Geosciences
Building #77
University of Arizona
Tucson, AZ 85721

Dr. William Wortman
Mission Research Corporation
735 State Street
P.O. Drawer 719
Santa Barbara, CA 93102

U.S. GOVERNMENT AGENCIES

Mr. Alfred Lieberman
ACDA/VI-OA, Room 5726
320 21st Street, N.W.
Washington, DC 20451

Colonel Jerry J. Perrizo
AFOSR/NP, Building 410
Bolling AFB
Washington, DC 20331-6448

Dr. Robert Blandford
AFTAC/CSS
1300 No. 17th St., Suite 1450
Arlington, VA 22209

AFTAC/CA
(STINFO)
Patrick AFB, FL 32925-6001

Dr. Frank F. Pilotte
HQ AFTAC/TT
Patrick AFB, FL 32925-6001

Katie Poley
CIA-ACIS/TMC
Room 4X16NHB
Washington, DC 20505

Dr. Larry Turnbull
CIA-OSWR/NED
Washington, DC 20505

Dr. Ralph W. Alewine, III
Dr. Alan S. Ryall, Jr.
Ms. Ann U. Kerr
DARPA/NMRO
1400 Wilson Blvd.
Arlington, VA 22209-2308
(7 copies)

DARPA/OASB/Librarian
1400 Wilson Blvd.
Arlington, VA 22209-2308

Dr. Dale Glover
DIA/DT-1B
Washington, DC 20301

Dr. Michael Shore
Defense Nuclear Agency/SPSS
6801 Telegraph Road
Alexandria, VA 22310

Dr. Max Koontz
U.S. Dept of Energy/DP-5
Forrestal Building
1000 Independence Avenue
Washington, DC 20585

Defense Technical Information Center
Cameron Station
Alexandria, VA 22314 (2 copies)

Dr. John J. Cipar, PL/LW
Phillips Lab/Geophysics Directorate
Hanscom AFB, MA 01731

James F. Lewkowicz, PL/LW
Phillips Lab/Geophysics Directorate
Hanscom AFB, MA 01731

Phillips Laboratory (PL/XO)
Hanscom AFB, MA 01731

Dr. James Hannon
Lawrence Livermore National Laboratory
P.O. Box 808
Livermore, CA 94550 (2 copies)

Office of the Secretary of Defense
DDR&E
Washington, DC 20330

Eric Chael
Division 9241
Sandia Laboratory
Albuquerque, NM 87185

Dr. William Leith
U.S. Geological Survey
Mail Stop 928
Reston, VA 22092

Dr. Robert Masse
Box 25046, Mail Stop 967
Denver Federal Center
Denver, CO 80225

Dr. Robert Reinke
WL/NTESG
Kirtland AFB, NM 87117-6008

CDRL MAILING LIST-NM

ORGANIZATION	NAME	NO. COPIES
NON-GOVERNMENT CONTRACTORS		
CALTECH	AHRENS	1
SAIC, SAN DIEGO	BACHE, SERENO	2
CANADA, GEOL SURVEY	BASHAM	1
ENSCO, SPRINGFIELD, VA	BAUMGARDT/DER	1
UCSD	BERGER	1
VPI	BOLLINGER	1
SAIC, ROSSLYN	BRATT, CARTER, LIBRARIAN	3
TELEDYNE, GARLAND, TX	BROWNE	1
WOODWARD-CLYDE	BURDICK	1
SHI	CHEFFY	1
U, TORONTO	CHUN	1
UCLA	DAVIS	1
SAN DIEGO STATE U.	DAY	1
SWEDEN, NAT. DEF. RES. INST.	EVA JOHANNISSON	1
MRC, SANTA BARBARA	FRISK	1
UCSC	FLATTE	1
PSR	FRITZEL	1
GERMANY, RUHR U	HARIES	1
CALTECH	HELMBERGER	1
SMUGEOPHYS. LAB	HERRIN, STUMP	1
CORNELL	ISACKS, BARAZANGI	1
UCB	JOHNSON, MCEVILLY	1
ENSCO, MELBOURNE, FL	KEMERAIT	1
ANU	KENNETT	1
LINCOLN LAB	LACOSS	1
U. ILL	LAMB	1
PENN STATE U.	LANGSTON	1
UCSC	LAY, SCHWARTZ	1
LDGO	LEPNER-LAM/RICHARDS	1
GERMANY, FED INST	MANFRED HENGER	1
ANFE	MARSHALL	1
NER	MARTIN	1
FRANCE, RADIOMANA	MASSINON, MECHELER	2
SMUPHYSICS DEPT	MCCARTOR, GRAY	1
S-CUBED, LA JOLLA	MCLAUGHLIN	1
UCSD	MINSTER, ORCUTT, GIVEN	2
ST LOUIS U	MITCHELL, HERFMANN	1
S-CUBED, PISTON	MURPHY	2
RADDX	PULLI	1
NORWAY, NTNF	RINGDAL	2
TELEDYNE, ALEXANDRIA, VA	RIVERS	2
TASC	SAILOR	1

CORL MAILING LIST-NM

ORGANIZATION	NAME	NO. COPIES
USC	SAMMIS, AKI	1
IRIS	SIMPSON	2
U. WASHINGTON	SMITH	1
U. WISCONSIN	THURBER, MEYER	1
MIT	TOKSOZ/DAINTY	1
U. AZ	WALLACE	1
MRC, NEWINGTON, VA	WORTMAN	1
US GOVERNMENT AGENCIES		
ACDA	LIEBERMAN	1
APOSRAP	JERRY PERRIZO	1
AFTAC, CSS, ROSSLYN, VA	BLANDFORD	1
AFTAC/CA	STINFO	1
AFTAC/TT	PILOTTE	1
CIA/ACIS	KATIE POLEY	1
CIA/OSWR	TURNBULL	1
DARPA	ALEWINE, RYALL, KERR	7
DARPA/RMO	LIBRARIAN	1
DIA	GLOVER	1
DNA/SPSS	SHORE	1
DOE	KOONTZ	1
DTIC	INFO CTR	2
GLLWH	CIPAR	1
GLLWH	LEWKOWICZ	1
GL/XO	XO	1
LLNL	HANNON	2
OSD	DOFE	1
SANDIA	CHAEI	1
USGS	LEITH	1
USGS	MASSE	1
WL/NTESG	REINKE	1
TOTAL NUMBER OF REPORTS		88